

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : C11D 3/386	A1	(11) International Publication Number: WO 88/ 09367 (43) International Publication Date: 1 December 1988 (01.12.88)
(21) International Application Number: PCT/US88/01844 (22) International Filing Date: 31 May 1988 (31.05.88) (31) Priority Application Number: 056,500 (32) Priority Date: 29 May 1987 (29.05.87) (33) Priority Country: US (60) Parent Application or Grant (63) Related by Continuation US 056,500 (CIP) Filed on 29 May 1987 (29.05.87) (71) Applicant (for all designated States except US): GEN- ENCOR, INC. [US/US]; 180 Kimball Way, So. San Francisco, CA 94080 (US).		(72) Inventors; and (75) Inventors/Applicants (for US only) : KOLATTUKUDY, Pappachan [US/US]; 2301 Hoxton Court, Columbus, OH 43220 (US). POULOSE, Ayrookaran, J. [IN/US]; 2540 Carmel Drive, San Bruno, CA 94066 (US). (74) Agent: PASSE, James, G.; Genencor, Inc., 180 Kimball Way, So. San Francisco, CA 94080 (US). (81) Designated States: AT (European patent), BE (Euro- pean patent), CH (European patent), DE (European patent), FR (European patent), GB (European pa- tent), IT (European patent), JP, LU (European pa- tent), NL (European patent), SE (European patent), US. Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt</i> <i>of amendments.</i>
(54) Title: CUTINASE CLEANING COMPOSITION (57) Abstract This invention relates to cleaning compositions and methods for using them. Particularly, the invention relates to compositions comprising a surfactant and a cutinase enzyme.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

Cutinase Cleaning Compositions

Background of the Invention

a) Field of the Invention

The invention relates to enzymatic cleaning compositions and methods for using them. Particularly the invention relates to cleaning compositions comprising a surfactant and a cutinase enzyme.

b) Background Information

A wide variety of enzymes are well known for use in cleaning compositions. The use of B. subtilis and B. licheniformis protease in commercial preparations is common. Other enzymes have also been used in commercial cleaning compositions such as, for example, U.S. Patent No. 4,011,169, and British Patent No. 1,293,613. Also a comprehensive review article of lipases in cleaning compositions can be found in Journal of Applied Biochemistry, 2:218-229 (1980) in an article entitled "Lipases as Detergent Components". Lipolytic detergent additives are also known from e.g., British Patent Specification No. 1,293,613 and Canadian Patent No. 835,343.

U.S. Patent No. 3,950,277 and British Patent Specification No. 1,442,418 disclose lipase enzymes combined with an activator and calcium and/or magnesium ions, respectively, which are utilized to pre-soak soiled fabrics and to remove triglyceride stains and soils from polyester or polyester/cotton fabric blends, respectively. Suitable microbial lipases for use therein (apart from animal and plant derived lipases) are said to be those derived from Pseudomonas, Aspergillus, Pneumococcus, Staphylococcus, and Staphylococcus toxins, Mycobacterium tuberculosis, Mycotorula lipolytica, and Sclerotinia.

British Patent Specification No. 1,372,034 discloses a detergent composition comprising a bacterial lipase produced by Pseudomonas stutzeri strain ATCC 19154. Furthermore, it is recommended that the preferred lipolytic enzymes should have a pH optimum between 6 and 10, and should be active in said range, preferably between 7 and 9. Around 1970, this presumed Pseudomonas stutzeri strain was reclassified as Pseudomonas aeruginosa, as appears for example from the ATCC catalogues.

European Patent Application EP-A-0130064 discloses an enzymatic detergent additive comprising a lipase isolated from Fusarium oxysporum with an alleged higher lipolytic cleaning efficiency than conventional lipases.

In European Patent Application No. 0214761, Enzymatic detergent additives are described as the active component of which is a microbially produced lipase from a strain of Pseudomonas cepacia. The lipases described therein are claimed to be superior to the lipolytic detergent action of the prior art, especially at low temperature washing processes (around 60°C and below).

In PCT Patent Application No. 87/00859 other novel lipolytic enzymes are described as having an optimal pH in the range of 8 - 10.5 at a temperature of 60°C or less from bacterial strains selected from Pseudomonas pseudoalcaligenes, P. stutzeri and Acinetobacter calcoaceticus. These enzymes are described as particularly effective at low temperatures; i.e., 40°C or lower and effective in both liquid and solid detergent compositions.

Also in U.S. Patent No. 3,950,277, it is described in general terms that lipases from Pseudomonas are suited as agents for removal of oily stains from fabrics, if used together with a special group of lipase activators. The art cited does not, however, cover cutinase enzymes from Pseudomonas or any other microbial source. However, prior art enzymes for use in cleaning compositions, while effective on many proteins and lipids, are not completely effective against all stains commonly found in laundry and other cleaning applications.

SUBSTITUTE SHEET

Further, many lipases are not stable at pH 8 - 11 where most cleaning compositions are used. Even further, most enzymes for use in cleaning compositions are not very stable, if at all, under oxidative conditions or in the presence of other enzymes such as proteases.

SUMMARY OF THE INVENTION

Accordingly it has been discovered that combinations of a surfactant and a substantially pure microbial cutinase enzyme are effective compositions for cleaning applications. The cutinase enzyme preparations possess activity at pH of from about 8 and 11, exhibit cleaning activity in aqueous solution at concentrations from about .05 mg/L to about 100 mg/L or more at temperatures from about 20°C to about 50°.

The enzymes are oxidatively stable and stable in the presence of other enzymes such as proteases. Even further, the cutinases show a synergistic effect when a plurality of surfactants are used with the cutinase.

The invention also relates to the improved process for enzymatically cleaning a material with an aqueous solution; the improvement comprising adding a substantially pure cutinase to the cleaning solution.

DETAILED DESCRIPTION OF THE INVENTION

Applicant has discovered that cutinase enzymes are useful when included in cleaning compositions. These compositions may take on a variety of forms such as for laundry cleaning, household and industrial cleaning, and the like. The cleaning compositions comprise combinations of known surfactants and a microbial cutinase enzyme which can be used to clean a wide variety of materials. The composition can be added to aqueous solution or solid powder, or formulated in an aqueous solution or solid powder and used according to conventional cleaning techniques.

SUBSTITUTE SHEET

Enzyme

Cutinases are well known in the art and are available from a wide variety of sources. See *Cutinases from Fungi and Pollen*, P.E. Kolattukudy, pg. 472 - 504, incorporated herein by reference, for discussion of cutinases useful in the practice of the invention. A preferred cutinase is that cutinase isolated in a substantially pure form from *Pseudomonas putida*, particularly the *P. putida*, ATCC 53552, described in copending U.S. patent application, Serial No. 932,959 filed November 19, 1986 and incorporated herein by reference, which enzyme therefrom has the following amino acid sequence:

SUBSTITUTE SHEET

5

1	ala	pro	leu	pro	asp	thr	pro	gly	ala	10	pro	phe	pro
										20	arg	ser	gly
	ala	val	ala	asn	phe	ala	arg	ser	gly		pro	tyr	thr
										30	gly	pro	ser
	thr	ser	ser	gln	ser	glu	gly	pro	ser		cys	arg	ile
										40	gly	gln	gly
	tyr	arg	pro	arg	asp	leu	gly	gln	gly		gly	val	arg
										50	thr	gly	ala
	his	pro	val	ile	leu	trp	gly	asn	gly				
										70	ser	his	trp
	gly	pro	ser	thr	tyr	ala	gly	leu	leu				
										80	val	ala	ala
	ala	ser	his	gly	phe	val	val	ala	ala		ala	glu	thr
										90	arg	glu	met
	ser	asn	ala	gly	thr	gly	arg	glu	met		leu	ala	cys
										100	thr	pro	tyr
	leu	asp	tyr	leu	val	arg	glu	asn	asp				
										110	gly	arg	val
	gly	thr	tyr	ser	gly	lys	leu	asn	thr				
										120	gly	arg	val
										130	gly	gly	ser
	gly	thr	ser	gly	his	ser	gln	gly	gly				
										140	thr	arg	thr
	ile	met	ala	gly	gln	asp	thr	arg	val		arg	thr	thr
										150	thr	leu	gly
	ala	pro	ile	gln	pro	tyr	thr	leu	gly		leu	gly	his
										160	arg	glu	met
	asp	ser	ala	ser	gln	arg	arg	gln	gln		gly	pro	met
										170	thr	arg	thr
	phe	leu	met	ser	gly	gly	gly	asp	thr		ile	ala	phe
										180	thr	arg	ala
	pro	tyr	leu	asn	ala	gln	pro	val	tyr		arg	arg	ala
										190	thr	arg	ala
	asn	val	pro	val	phe	trp	gly	glu	arg		arg	tyr	val
										200	thr	arg	ala
										210	thr	arg	ala
	ser	his	phe	glu	pro	val	gly	ser	gly		gly	ala	tyr
										220	thr	arg	ala
	arg	gly	pro	ser	thr	ala	trp	phe	arg		phe	gln	leu
										230	thr	arg	ala
	met	asp	asp	gln	asp	ala	arg	ala	thr		phe	tyr	gly
										240	thr	arg	ala
	ala	gln	cys	ser	leu	cys	thr	ser	leu		leu	trp	
										250	thr	arg	ala
	ser	val	gly	arg	arg	gly	leu						

SUBSTITUTE SHEET

Other sources of bacterial and fungal cutinases include:

Fusarium solani pisi

Fusarium roseum sambucinum

Fusarium roseum culmorum

Helminthosporium sativum

Ulocladium consortiale

Streptomyces scabies

Colletotrichum capsici

Phytophthora cactorum

Botrytis cineria

Colletotrichum gloeosporioides

The cutinase of the invention should preferably be selected to cause at least about 10%, and preferably 20%, hydrolysis of the given fat under given conditions. Normally the amount would be in a concentration of from about .01% to about 5.0% by weight of the surfactant, and preferably from about .05% to about 3%, such that upon dilution in wash water it is in a concentration of at least about .05mg/L. Further, one skilled in the art could take the preferred cutinase or, for that matter, any cutinase of the invention or any immunologically identical cutinase and use random or selective replacement of amino acids to produce other cutinases which are more or less selective toward given substrates or include modification in activity such as oxidative stability. Substantially pure cutinase includes the isolated enzyme as well as the broth containing the enzyme in unpurified form but essentially free of other enzymes and enzyme sources.

The natural substrate of cutinase is cutin which is a biopolyester polymer which covers the plant leaves, fruits, etc., see Structure, Biosynthesis and Biodegradation of Cutin and Suberin. (1981), P.E. Kolattukudy, Ann. Rev. Plant Physiol., 32, pgs. 539-567. Stains comprising lipids which could be hydrolyzed or bound by cutinase on a substrate such as cloth would be similar to the natural substrate cutin. Cutinase, for these types of stains, will be more

SUBSTITUTE SHEET

effective than the prior art lipases. The cutinases will work especially well on gravy, oils and greases, plant or grass, oil based makeup and collar stains.

Cutinases are distinguishable from other lipases by methods well known in the art, See R.E. Purdy and P.E. Kolattukudy, Biochemistry, "Cutinase Assay", 14:2831-2840, (1975). Microbial cutinases from both fungal and bacterial sources have very good activity at pH 8 - pH 11 which is an ideal pH condition for detergent use.

Because of the specific activity of cutinases, it is a preferred aspect of the invention to combine one or more other cutinases, or one or more other enzymes, such as proteases, amylases or other lipases, along with the cutinase of the invention in the cleaning composition. Further, Applicant shows a synergistic increase in hydrolytic activity of cutinase when two or more surfactants are combined along with the cutinase enzyme.

Cutinases then are ideal for cleaning composition inclusion. They have stability oxidatively such as in H_2O_2 . They have good stability in a temperature range of from about 20 - 50°C which is ideal from a cleaning point of view. They are also stable in the presence of other enzymes; e.g., proteases, and as such, are ideal for mixtures of enzymes.

The Surfactant

A number of known compounds are suitable surfactants useful in the present compositions. These include nonionic, anionic, cationic, or zwitterionic detergents, as disclosed in U.S. 4,404,128 to Barry J. Anderson and U.S. 4,261,868 to Jiri Hora et al. The art is familiar with the different formulations which can be used as cleaning compositions.

SUBSTITUTE SHEET

The Cleaning Compositions and Method of Use

Cutinases can be formulated as a purposefully added ingredient into known powdered and liquid detergents having pH between 6.5 and 12.0 at levels of about 0.01 to about 5% (preferably .05 to .5%) by weight of the detergent. These detergent cleaning compositions can also include other enzymes such as known proteases and amylases, as well as bleaches, colorants, builders, and stabilizers.

The cutinase of the invention may be added to powdered detergents in the form of granulates or prills, prepared by methods known in the art such as described in British Patent Nos. 1,324,116 and 1,362,365 and U.S. patent Nos. 3,519,570; 4,106,991 and 4,242,219.

The cutinase preparations of the invention can be prepared by cultivating the microorganisms defined herein or otherwise cutinase containing microorganism under appropriate conditions. In order to obtain reasonable yields of enzyme, media containing readily assimilable carbon and energy sources as necessary such as a nitrogen source, as well as calcium and magnesium salts and trace elements and cutin, or monomers of cutin, or compounds resembling cutin or cutin monomers. One could also obtain the gene for cutinase and express in any organism of choice where one may not have to add cutin or cutin monomers into the fermentation.

The addition of cutinase to conventional cleaning compositions does not create any special use limitation. In other words, any temperature and pH suitable for detergent compositions containing enzymes is also suitable for the present compositions.

Although the preferred form of the invention has been described above, it will be obvious to those skilled in the art to which the invention pertains, that, after understanding the invention and in view of the following testing as a whole, various changes and equivalent modifications may be made without parting from the scope of the invention as defined by the claims.

SUBSTITUTE SHEET

STABILITY CUTINASE AGAINST PROTEASES

Reaction conditions:

Buffer: 0.1 M NaP, pH 10

Temp: 37°C

Lipase: 42 ug/ml

Approximately 1:1 protease:cutinase aqueous solution were made up with the following results.

ENZYME ACTIVITY

Protease

Incubation Time	0 min	5 min	10 min	15 min	14 hrs
None	1.57	1.60	1.47	1.63	1.61
Maxacal (35ug/ml)	1.68	1.58	1.72	1.66	0.134
Esperase (64 ug/ml)	1.73	1.64	1.59	1.51	0.456

Maxical is Gist-Brocade's brand of subtilisin enzyme (protease)

Esperase is Novo's brand of protease enzyme (protease)

SUBSTITUTE SHEET

TEMPERATURE STABILITY OF BACTERIAL CUTINASE

HALF LIFE AT 50°C

<u>pH</u>	<u>Hrs.</u>
7	30
8	25
9	12
10	0.3

Enzyme was incubated at 50°C in 0.1M sodium phosphate buffer at various pH's and activity was measured by hydrolysis of trioctanoin in polyvinyl alcohol emulsions.

SUBSTITUTE SHEET

EFFECT OF DETERGENTS ON HYDROLASE ACTIVITY

Reaction Conditions:

substrate: p-nitro-phenyl butyrate, 1mM (pnp)

pH: 8.0

buffer: 0.1M tris pH 8.0

temperature: 25°C

enzyme: bacterial cutinase from ATCC 53552

An aqueous solution with the following were made up and the enzyme activity was measured in these solutions using pnp as a substrate by following absorbance of p-nitrophenol at 410 nm.

<u>Triton x100%</u>	<u>SDS %</u>	<u>% Activity</u>
0	0	100
0.2	—	78
0.4	—	60
—	0.05	30
—	0.1	23
—	0.2	14
—	0.4	6
0.4	0.4	78
0.2	0.2	98
0.2	0.05	125
0.2	0.1	138

SUBSTITUTE SHEET

6.1	0.1	130
0.05	0.05	132

- 1) Non-ionic detergent inhibition is not significant at low concentrations.
- 2) An ionic detergent inhibitor at high concentrations.
- 3) Mixture of anionic and non-ionic detergents stimulate activity.

SUBSTITUTE SHEET

STABILITY TOWARD OXIDANTS

Cutinase .5 mg/ml in 0.1M sodium phosphate buffer, was incubated with various levels of hydrogen peroxide at pH 8.4 and 25°C for 2 hours, and hydrolytic activity was measured by a pH-stat using trioctanoim-polyvinyl alcohol emulsion.


[H ₂ O ₂ ppm]	<u>Hydrolytic Activity</u>
	<u>% Remaining</u>
0	100
100	86
200	86
500	91
1000	95

SUBSTITUTE SHEET

TABLE 1

pH—optimum of bacterial cutinase.

50 mM NaP, 30 C, 0.5 % TO.




SUBSTITUTE SHEET

TABLE 2

Temperature optimum of bact. cutinase.

50 mM NaP, pH 8.0.



SUBSTITUTE SHEET

What is Claimed is:

1. An enzymatic cleaning composition comprising a substantially pure microbial cutinase in an amount sufficient that upon dilution in an aqueous solution for use there is sufficient cutinase to achieve a concentration of from about .05 mg/L to about 100 mg/L of cutinase.
2. A composition according to Claim 1 wherein the cutinase is the cutinase present in an amount of from about 0.01% to about 5% by weight of the surfactant.
3. A composition according to Claim 2 wherein the Pseudomonas putida is ATCC 53552.
4. A composition according to Claim 1 which further comprises one or more enzymes selected from lipases, amylases or proteases.
5. A composition according to Claim 1 wherein the cutinase selected is capable of causing at least 10% hydrolysis of a cutin or cutin-like substrate desirous of being cleaned from a given surface.
6. A composition according to Claim 1 wherein the cutinase is a non-naturally occurring cutinase having at least one amino acid randomly or selectively replaced by an amino acid not naturally found at that position.

SUBSTITUTE SHEET

7. A composition according to Claim 1 wherein the cutinase is added in the form of cell fermentation broth containing cutinase but essentially free of other enzymes or enzyme sources.
8. An enzymatic cleaning additive wherein the active component which is a substantially pure microbially produced cutinase.
9. A composition according to Claim 8 which further comprises one or more enzymes selected from lipases, amylases or proteases.
10. A composition according to Claim 8 comprising a plurality of surfactants.
11. A composition according to Claim 10 wherein the surfactants are SDS and Triton X-100.
12. An improved method for enzymatically cleaning a material having a cutin or cutin-like stain comprising:
 - a) selecting a cutinase enzyme;
 - b) forming an aqueous solution with said enzyme in a concentration of from about 0.05 mg/L to about 100 mg/L;
 - c) contacting the material with the solution of step b); and
 - d) rinsing the material of step c).
13. A method according to Claim 12 wherein the aqueous solution further comprises a surfactant compatible with the selected cutinase.

14. A method according to Claim 13 wherein the aqueous solution further comprises a surfactant compatible with the selected cutinase.

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 88/01844


I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶
 According to International Patent Classification (IPC) or to both National Classification and IPC
 IPC⁴: C 11 D 3/386

II. FIELDS SEARCHED
 Minimum Documentation Searched ⁷
 Classification System Classification Symbols
 IPC⁴ C 11 D; C 12 N
 Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸

III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
P,X	EP, A, 0268452 (GENENCOR) 25 May 1988 see example 13; abstract; page 1, lines 29-30 cited in the application -----	1-3,5,6,8, 12,13

- ¹⁰ Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
 - "E" earlier document but published on or after the international filing date
 - "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 - "O" document referring to an oral disclosure, use, exhibition or other means
 - "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "G" document member of the same patent family

IV. CERTIFICATION	
Date of the Actual Completion of the International Search 29th August 1988	Date of Mailing of this International Search Report 20 SEP 1988
International Searching Authority EUROPEAN PATENT OFFICE	Signature of Authorized Officer  P.C.G. VAN DER PUTTEN

SA 22728

6450 FORM 10479

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0268452	25-05-88	AU-A- - 8115387 AU-A- 7901387	26-05-88 09-06-88
<hr/>			